

Index and secondary index theory for flat bundles with duality

Ulrich Bunke*and Xiaonan Ma[†]

February 1, 2008

Abstract

We discuss some aspects of index and secondary index theory for flat bundles with duality. This theory was first developed by Lott [9]. Our main purpose in the present paper is provide a modification with better functorial properties.

Contents

1	The functor L	1
1.1	Definition of L	1
1.2	Computation of $L(X)$	1
1.3	The natural transformation to K -theory	2
1.4	Secondary L -groups. The $\mathbb{R}/\mathbb{Z}$ -variant	3
1.5	Index and secondary index	3
2	Geometry and extended secondary L-groups	5
2.1	Definition of $\bar{L}$	5
2.2	Functorial properties	6
3	Eta invariants and index maps	8
3.1	The η -invariant	8
3.2	The secondary index map	9
3.3	The index map for L^{ex} and $\bar{L}^{ex}$	10

*Göttingen, bunke@uni-math.gwdg.de

[†]Berlin, ma@mathematik.hu-berlin.de

1 The functor L

1.1 Definition of L

We define a contravariant functor L from the category $\mathcal{Top}$ of topological spaces and continuous maps to $\mathbb{Z}_2$ -graded rings.

Let X be a topological space and $\epsilon \in \mathbb{Z}_2$. If $\mathcal{F}$ is a locally constant sheaf of finite-dimensional $\mathbb{R}$ -modules over X , then an ϵ -**symmetric duality structure** on $\mathcal{F}$ is an isomorphism of sheafs $q : \mathcal{F} \rightarrow \mathcal{F}^*$ satisfying $q^* = (-1)^\epsilon q$, where $\mathcal{F}^* := \text{Hom}(\mathcal{F}, \underline{\mathbb{R}})$.

We first define a semigroup $\hat{L}_\epsilon(X)$ and then obtain $L_\epsilon(X)$ by introducing a relation. A generator of the semigroup $\hat{L}_\epsilon(X)$ is a pair $(\mathcal{F}, q)$ consisting of a locally constant sheaf of finite-dimensional $\mathbb{R}$ -modules and an ϵ -symmetric duality structure q . The operation in $\hat{L}_\epsilon(X)$ is given by direct sum

$$(\mathcal{F}, q) + (\mathcal{F}', q') := (\mathcal{F} \oplus \mathcal{F}', q \oplus q') .$$

The group $L_\epsilon(X)$ is obtained from $\hat{L}_\epsilon(X)$ by introducing the relation **lagrangian reduction**. We require $(\mathcal{F}, q) \sim 0$ if there exists a locally constant lagrangian subsheaf $i : \mathcal{L} \hookrightarrow \mathcal{F}$, i.e. the composition $\mathcal{L} \xrightarrow{i} \mathcal{F} \xrightarrow{q} \mathcal{F}^* \xrightarrow{i^*} \mathcal{L}^*$ vanishes and $\mathcal{L} = \mathcal{L}^\perp := \ker(i^* \circ q)$.

The class of $(\mathcal{F}, q)$ in $L_\epsilon(X)$ will be denoted by $[\mathcal{F}, q]$. The ring structure $L_\epsilon(X) \otimes L_{\epsilon'}(X) \rightarrow L_{\epsilon+\epsilon'}(X)$ is induced by the tensor product:

$$[\mathcal{F}, q][\mathcal{F}', q'] := \frac{\sqrt{(-1)^\epsilon} \sqrt{(-1)^{\epsilon'}}}{\sqrt{(-1)^{\epsilon+\epsilon'}}} [\mathcal{F} \otimes \mathcal{F}', q \otimes q'] .$$

The sign-convention is made such that later we have a natural transformation of rings from L to complex K -theory K^0 .

If $f : Y \rightarrow X$ is a morphism in $\mathcal{Top}$, then $f^* : L(X) \rightarrow L(Y)$ is defined by $f^*[\mathcal{F}, q] = [f^*\mathcal{F}, f^*q]$. The map f^* only depends on the homotopy class of f .

Remark: A version $L(X)^{\text{Lott}}$ of this ring was first introduced by Lott [9]. His definition differs from ours since our relation "lagrangian reduction" is replaced by "hyperbolic reduction" in the definition of Lott. Here a generator $(\mathcal{F}, q)$ is called hyperbolic if there is a lagrangian subsheaf $\mathcal{L} \subset \mathcal{F}$ which admits a complement inside $\mathcal{F}$. In particular, $L(X)$ is a quotient of $L(X)^{\text{Lott}}$.

1.2 Computation of $L(X)$

Assume that X is path connected and fix a base point x . Let $G := \pi_1(X, x)$ be the fundamental group. We consider the semigroup $\hat{L}_\epsilon(G)$ of all tuples (F, q, ρ) , where F is a finite-dimensional real vector space, $q : F \rightarrow F^*$ is an ϵ -symmetric duality structure, and $\rho : G \rightarrow \text{Aut}(F, q)$ is a representation of G . The semigroup operation is given by direct sum. We then define the

group $L_\epsilon(G)$ by introducing the relation $(F, q, \rho) \sim 0$ if there exists an invariant lagrangian subspace $L \subset F$. Taking the holonomy representation on the fibre over x we obtain a bijection $L(X) \cong L(G)$.

Let $\hat{G} := G \times \mathbb{Z}_2$. If $(F, q, \rho) \in \hat{L}(G)$, then we form the representation $(\hat{F}, \hat{\rho})$ of $\hat{G}$ by $\hat{F} := F \oplus F^*$, $\hat{\rho}((g, 0)) := \rho(g) \oplus \rho(g^{-1})^*$,

$$\hat{\rho}(1, 1) := \begin{pmatrix} 0 & q^{-1} \\ q & 0 \end{pmatrix}.$$

If (F, q, ρ) is an irreducible representation of G , then $(\hat{F}, \hat{\rho})$ is an irreducible representation of $\hat{G}$. An irreducible representation (V, σ) of $\hat{G}$ is called real, complex, or quaternionic, if $\text{End}_G(V) \cong \mathbb{R}$, $\cong \mathbb{C}$ or $\cong \mathbb{H}$, respectively.

For each irreducible representation (F, ρ) of G we define a $\mathbb{Z}_2$ -graded group $A(F, \rho)$ as follows. If (F, ρ) admits an invariant ϵ -symmetric form q , then we define $A_\epsilon(F, \rho) := \mathbb{Z}$, iff $(\hat{F}, \hat{\rho})$ is real or quaternionic, $A_\epsilon(F, \rho) := \mathbb{Z}_2$, iff $(\hat{F}, \hat{\rho})$ is complex, and we set $A_\epsilon(F, \rho) := 0$, if (F, ρ) does not admit an invariant ϵ -symmetric form. If $A(F, \rho) \neq 0$, then we fix an invariant ϵ -symmetric form $q_{F, \rho}$.

Let $\text{Rep}(G)$ denote the set of isomorphism classes of irreducible representations of G .

Theorem 1.1 *There is a natural isomorphism of $\mathbb{Z}_2$ -graded groups*

$$L(G) = \bigoplus_{(F, \rho) \in \text{Rep}(G)} A(F, \rho)$$

fixed by the condition that $(F, q, q_{F, q}) \mapsto 1_{(F, q)}$.

1.3 The natural transformation to K -theory

By $\mathcal{T}op_{para}$ we denote the full subcategory of $\mathcal{T}op$ of paracompact topological spaces. Let $K^0(X)$ be the complex K -theory functor. We construct a natural transformation $b : L \rightarrow K^0$ of functors from $\mathcal{T}op_{para}$ to rings.

A locally constant sheaf of finite-dimensional $\mathbb{R}$ -modules on X gives rise to a locally trivial real vector bundle $bundle(\mathcal{F})$ in a natural way. The correspondence $bundle$ is functorial and compatible with direct sum, tensor product, and duality. Thus applying the bundle construction to $(\mathcal{F}, q)$ we obtain a pair (F, Q) consisting of a finite-dimensional real vector bundle and an isomorphism $Q : F \rightarrow F^*$.

Let (F, Q) be a real vector bundle with an isomorphism $Q : F \rightarrow F^*$ such that $Q^* = (-1)^\epsilon Q$ for $\epsilon \in \mathbb{Z}_2$. Following the language introduced by Lott [9] we call an isomorphism $J : F \rightarrow F$ a metric structure, if

1. $J^* \circ Q$ defines a scalar product on F ,

$$2. J^2 = (-1)^\epsilon \text{id}_F,$$

$$3. J^* \circ Q = (-1)^\epsilon Q \circ J.$$

Since we assume that X is paracompact it admits partitions of unity. This implies that metric structures exist, and that the space of all metric structures is contractible.

Given (F, Q) as above we choose a metric structure J . Let $F_{\mathbb{C}}$ be the complexification of F . Then $\frac{1}{\sqrt{(-1)^\epsilon}}J$ is a $\mathbb{Z}_2$ -grading of $F_{\mathbb{C}}$, and the pair $(F_{\mathbb{C}}, \frac{1}{\sqrt{(-1)^\epsilon}}J)$ represents an element of $K^0(X)$ which does not depend on the choice of J .

The transformation $b : L \rightarrow K^0$ is obtained by composing the latter construction with *bundle*.

1.4 Secondary L -groups. The $\mathbb{R}/\mathbb{Z}$ -variant

We first recall the definition of the 2-periodic cohomology theory $K_{\mathbb{R}/\mathbb{Z}}$ introduced by [7], [8]. Let BU be the classifying space of complex K -theory. The Chern character (with real coefficients) is induced by a map $\mathbf{ch} : BU \rightarrow \prod_{n=1}^{\infty} K(\mathbb{R}, 2n)$. The homotopy fibre of this map classifies $K_{\mathbb{R}/\mathbb{Z}}$. In particular, for any paracompact space X there is a natural exact sequence of $K^0(X)$ -modules

$$\rightarrow K^{-1}(X) \xrightarrow{\mathbf{ch}} H^{odd}(X, \mathbb{R}) \rightarrow K_{\mathbb{R}/\mathbb{Z}}^{-1}(X) \xrightarrow{\beta} K^0(X) \xrightarrow{\mathbf{ch}} H^{ev}(X, \mathbb{R}) \rightarrow ,$$

where $K^0(X)$ acts on cohomology via the Chern character.

We now define the functor $X \mapsto \bar{L}^{\mathbb{R}/\mathbb{Z}}(X)$ from paracompact topological spaces to $\mathbb{Z}_2$ -graded groups by the following pull-back diagram:

$$\begin{array}{ccc} \bar{L}^{\mathbb{R}/\mathbb{Z}}(X) & \rightarrow & L(X) \\ \downarrow & & \downarrow b \\ K_{\mathbb{R}/\mathbb{Z}}^{-1}(X) & \xrightarrow{\beta} & K^0(X) \end{array} .$$

The grading is induced from that of $L(X)$. On morphisms the functor $L^{\mathbb{R}/\mathbb{Z}}$ only depends on homotopy classes. Note that $K^0(X)$ and $K_{\mathbb{R}/\mathbb{Z}}^{-1}(X)$ are $L(X)$ -modules via b . This induces a graded $L(X)$ -module structure on $\bar{L}^{\mathbb{R}/\mathbb{Z}}(X)$. We have the following natural exact sequence of $L(X)$ -modules

$$K^{-1}(X) \xrightarrow{\mathbf{ch}} H^{odd}(X, \mathbb{R}) \rightarrow \bar{L}^{\mathbb{R}/\mathbb{Z}}(X) \rightarrow L(X) \xrightarrow{\mathbf{ch} \circ b} H^{ev}(X, \mathbb{R}) .$$

1.5 Index and secondary index

Let $X \rightarrow B$ be a smooth locally trivial fibre bundle over a compact base B with compact even-dimensional fibres such that the vertical bundle TX/B is orientable. In this case we have the following maps:

- $\pi_*^{sign} : H^*(X, \mathbb{R}) \rightarrow H^*(B, \mathbb{R})$ defined by $\pi_*(\omega) = \int_{X/B} \omega \cap \mathbf{L}(TX/B)$, where $\int_{X/B}$ is integration over the fibre and $\mathbf{L}(TX/B)$ denotes the Hirzebruch $\mathbf{L}$ -class of the vertical bundle.
- $\pi_!^{sign} : K^0(X) \rightarrow K^0(B)$ defined by $\pi_!^{sign}([E]) = \text{index}(D_E^{sign})$, where D_E^{sign} is the fibrewise signature operator twisted by E and $\text{index}(D_E^{sign}) \in K^0(B)$ denotes the class of the index bundle.
- $\pi_*^L : L(X) \rightarrow L(B)$ defined by $\pi_*^L[\mathcal{F}, q] = [H^\bullet R\pi_* \mathcal{F}, \pi_*(q)]$, where $H^\bullet R\pi_* \mathcal{F} = \bigoplus_{i=0}^\infty R^i \pi_*(\mathcal{F})$ is the direct sum of higher derived direct images of $\mathcal{F}$. The orientation of TX/B provides an isomorphism $p : R^{\dim(TX/B)} \pi_* \mathbb{R} \xrightarrow{\sim} \mathbb{R}$. It induces isomorphisms $R^i \pi_* \mathcal{F}^* \rightarrow (R^{\dim(TX/B)-i} \pi_* \mathcal{F})^*$, and $\pi_*(q)$ is the composition of the sum of these maps with the sum of $R^i \pi_*(q) : R^i \pi_* \mathcal{F} \rightarrow R^i \pi_* \mathcal{F}^*$.

It is an exercise in spectral sequences to show that π_*^L is well-defined.

Remark: In [9] Lott defines $\pi_*^{L, Lott} : L^{Lott}(X) \rightarrow L^{Lott}(B)$ which induces π_*^L by passing to quotients.

By the index theorem for families and fibrewise Hodge theory the following diagram commutes

$$\begin{array}{ccccc} L(X) & \rightarrow & K^0(X) & \rightarrow & H^{ev}(X, \mathbb{R}) \\ \pi_*^L \downarrow & & \pi_!^{sign} \downarrow & & \pi_*^{sign} \downarrow \\ L(B) & \rightarrow & K^0(B) & \rightarrow & H^{ev}(B, \mathbb{R}) \end{array} .$$

In order to define an index map for $K_{\mathbb{R}/\mathbb{Z}}$ we need the further assumption that π is K -oriented. Thus assume that TX/B has a $Spin_c$ -structure. Then there are maps $\pi_!^{Spin_c} : K^0(X) \rightarrow K^0(B)$ and $\pi_!^{Spin_c, \mathbb{R}/\mathbb{Z}} : K_{\mathbb{R}/\mathbb{Z}}^{-1}(X) \rightarrow K_{\mathbb{R}/\mathbb{Z}}^{-1}(B)$ (compare e.g. [8]), such that the following diagram commutes:

$$\begin{array}{ccccccc} H^{odd}(X, \mathbb{R}) & \rightarrow & K_{\mathbb{R}/\mathbb{Z}}^{-1}(X) & \rightarrow & K^0(X) & \rightarrow & H^{ev}(X, \mathbb{R}) \\ \pi_*^{Spin_c} \downarrow & & \pi_!^{Spin_c, \mathbb{R}/\mathbb{Z}} \downarrow & & \pi_!^{Spin_c} \downarrow & & \pi_*^{Spin_c} \downarrow \\ H^{odd}(B, \mathbb{R}) & \rightarrow & K_{\mathbb{R}/\mathbb{Z}}^{-1}(B) & \rightarrow & K^0(B) & \rightarrow & H^{ev}(B, \mathbb{R}) \end{array} ,$$

where $\pi_*^{Spin_c}(\omega) = \int_{X/B} \hat{\mathbf{A}}(TX/B) \cap e^{c_1/2} \cap \omega$ and c_1 is the first Chern class determined by the $Spin_c$ -structure. There is a unique element $E_{sign} \in K_0(X)$ such that $\pi_!^{sign}(x) = \pi_!^{Spin_c}(E_{sign} \bullet x)$. Note that $\mathbf{ch}(E_{sign}) \cap \hat{\mathbf{A}}(TX/B) \cap e^{c_1/2} = \mathbf{L}(TX/B)$.

- $\pi_*^{sign, \mathbb{R}/\mathbb{Z}} : K_{\mathbb{R}/\mathbb{Z}}^{-1}(X) \rightarrow K_{\mathbb{R}/\mathbb{Z}}^{-1}(B)$ is defined by $\pi_*^{sign, \mathbb{R}/\mathbb{Z}}(x) := \pi_*^{Spin_c, \mathbb{R}/\mathbb{Z}}(E_{sign} \bullet x)$ so that the following diagram commutes

$$\begin{array}{ccccccc} H^{odd}(X, \mathbb{R}) & \rightarrow & K_{\mathbb{R}/\mathbb{Z}}^{-1}(X) & \rightarrow & K^0(X) & \rightarrow & H^{ev}(X, \mathbb{R}) \\ \pi_*^{sign} \downarrow & & \pi_!^{sign, \mathbb{R}/\mathbb{Z}} \downarrow & & \pi_!^{sign} \downarrow & & \pi_*^{sign} \downarrow \\ H^{odd}(B, \mathbb{R}) & \rightarrow & K_{\mathbb{R}/\mathbb{Z}}^{-1}(B) & \rightarrow & K^0(B) & \rightarrow & H^{ev}(B, \mathbb{R}) \end{array} .$$

- $\pi_*^{\bar{L}, \mathbb{R}/\mathbb{Z}} : \bar{L}^{\mathbb{R}/\mathbb{Z}}(X) \rightarrow \bar{L}^{\mathbb{R}/\mathbb{Z}}(B)$ is the map induced by $\pi_!^{sign, \mathbb{R}/\mathbb{Z}}$ and π_*^L . The following diagram commutes:

$$\begin{array}{ccccccc}
H^{odd}(X, \mathbb{R}) & \rightarrow & \bar{L}^{\mathbb{R}/\mathbb{Z}}(X) & \rightarrow & L(X) & \rightarrow & H^{ev}(X, \mathbb{R}) \\
\pi_*^{sign} \downarrow & & \pi_*^{\bar{L}, \mathbb{R}/\mathbb{Z}} \downarrow & & \pi_*^L \downarrow & & \pi_*^{sign} \downarrow \\
H^{odd}(B, \mathbb{R}) & \rightarrow & \bar{L}^{\mathbb{R}/\mathbb{Z}}(B) & \rightarrow & L(B) & \rightarrow & H^{ev}(B, \mathbb{R})
\end{array}$$

All index maps are natural with respect to pull-back of fibre bundles.

Let $\pi_1 : X \rightarrow X_1$ and $\pi_2 : X_1 \rightarrow B$ be locally trivial smooth fibre bundles with closed even-dimensional fibres and compact base. Further assume that the vertical bundles TX_1/X_2 and TX_2/B carry $Spin_c$ -structures (and are therefore oriented). Then the composition $\pi = \pi_2 \circ \pi_1 : X \rightarrow B$ is a locally trivial fibre bundle with closed even-dimensional fibres, and the vertical bundle TX/B carries an induced $Spin_c$ -structure. In this situation the index maps on complex K -theory, $K_{\mathbb{R}/\mathbb{Z}}$ -theory, and in cohomology behave functorially with respect to the iterated fibre bundle. It is again an exercise in spectral sequences to show that π_*^L is also functorial. As a consequence, $\pi_*^{\bar{L}, \mathbb{R}/\mathbb{Z}}$ is functorial, i.e. $\pi_*^{\bar{L}, \mathbb{R}/\mathbb{Z}} = (\pi_2^{\bar{L}, \mathbb{R}/\mathbb{Z}})_* \circ (\pi_1^{\bar{L}, \mathbb{R}/\mathbb{Z}})_*$.

Remark : The motivation of introducing the quotient $L(X)$ of Lott's group $L^{Lott}(X)$ is to implement this functoriality for π_*^L , which is not true for Lott's definition.

2 Geometry and extended secondary L -groups

2.1 Definition of $\bar{L}$

The functor $X \mapsto \bar{L}^{\mathbb{R}/\mathbb{Z}}(X)$ from $\mathcal{Top}_{para}$ to $\mathbb{Z}_2$ -graded $L(X)$ -modules was defined by a purely homotopy-theoretic construction as an extension of the functor

$$X \mapsto \ker(\mathbf{ch} \circ b : L(X) \rightarrow H^{ev}(X, \mathbb{R}))$$

by $X \mapsto H^{odd}(X, \mathbb{R})/\mathbf{ch}(K^{-1}(X))$.

Let $\mathcal{Top}_{smooth}$ denote the full subcategory of topological spaces $\mathcal{Top}_{para}$ which are homotopy equivalent to smooth manifolds. In the present section we use a differential geometric construction in order to define on $\mathcal{Top}_{smooth}$ a functor $X \mapsto \bar{L}(X)$ to graded $L(X)$ -modules which extends $X \mapsto \ker(\mathbf{ch} \circ b)$ by $X \mapsto H^{odd}(X, \mathbb{R})$. First we define $\bar{L}(M)$ for a smooth manifold M . Again we first define a semigroup $\hat{\bar{L}}(M)$. A generator of $\hat{\bar{L}}(M)$ is a tuple $(\mathcal{F}, q, J, \rho)$, where $(\mathcal{F}, q)$ is a generator of $\hat{L}_\epsilon(M)$, J is a metric structure on $F := bundle(\mathcal{F})$, and $\rho \in \Omega^{4*-(-1)^\epsilon}(M)/\text{im}(d)$ satisfies $d\rho = p(\nabla^F, J)$. Here $(\Omega^*(M), d)$ is the real deRham complex of M and ∇^F is the canonical flat connection on F (such that $\ker(\nabla^F) \cong \mathcal{F}$). The metric structure J induces a $\mathbb{Z}_2$ -grading of $F_{\mathbb{C}}$, and if $\nabla^{F_{\mathbb{C}}, J}$ denotes the even part of the extension of ∇^F to

$F_{\mathbb{C}}$, then $p(\nabla^F, J) := \mathbf{ch}(\nabla^{F_{\mathbb{C}}, J}) = \text{tr}_s \exp(-R^{\nabla^{F_{\mathbb{C}}, J}}/2\pi i)$ is the characteristic form representing $\mathbf{ch} \circ b([\mathcal{F}, q])$. The semigroup operation is given by

$$(\mathcal{F}, q, J, \rho) + (\mathcal{F}', q', J', \rho') := (\mathcal{F} \oplus \mathcal{F}', q \oplus q', J \oplus J', \rho + \rho') .$$

We form $\bar{L}_\epsilon(M)$ by introducing the relation **lagrangian reduction**. Let $\mathcal{L} \subset \mathcal{F}$ be a locally constant lagrangian subsheaf and $L := \text{bundle}(\mathcal{L})$. Then we have a decomposition $F = L \oplus J(L)$. Let $\nabla^\oplus$ be the part of $\nabla^{F_{\mathbb{C}}, J}$ which preserves this decomposition, and let $\tilde{\mathbf{ch}}(\nabla^\oplus, \nabla^{F_{\mathbb{C}}, J})$ be the transgression Chern form such that $d\tilde{\mathbf{ch}}(\nabla^\oplus, \nabla^{F_{\mathbb{C}}}) = \mathbf{ch}(\nabla^\oplus) - \mathbf{ch}(\nabla^{F_{\mathbb{C}}})$. Note that $\mathbf{ch}(\nabla^\oplus) = 0$. In $\bar{L}_\epsilon(M)$ we require

$$(\mathcal{F}, q, J, \rho) \sim (0, 0, 0, \rho + \tilde{\mathbf{ch}}(\nabla^\oplus, \nabla^{F_{\mathbb{C}}, J})) .$$

By $[\mathcal{F}, q, J, \rho]$ we denote the class in $\bar{L}$ represented by $(\mathcal{F}, q, J, \rho)$.

Remark: In [9] similar functors $\bar{L}^{\text{Lott}}$ were defined replacing the relation "lagrangian reduction" by "hyperbolic reduction" and "change of metric structure". Lott's relation is smaller than ours, and we have a natural surjective map $\bar{L}^{\text{Lott}}(M) \rightarrow \bar{L}(M)$.

The graded module structure of $\bar{L}(X)$ over $L(X)$ is defined by

$$[\mathcal{F}, q, J, \rho] \bullet [\mathcal{E}, p] := \frac{\sqrt{(-1)^\epsilon} \sqrt{(-1)^{\epsilon'}}}{\sqrt{(-1)^{\epsilon+\epsilon'}}} [\mathcal{F} \otimes \mathcal{E}, q \otimes p, J \otimes J^E, \rho \wedge \mathbf{ch}(\nabla^{E_{\mathbb{C}}, J^E})] ,$$

where J^E is any metric structure on $\text{bundle}(\mathcal{E})$.

2.2 Functorial properties

If $f : M \rightarrow N$ is a smooth map of manifolds, then we obtain an induced map $f^* : \bar{L}(N) \rightarrow \bar{L}(M)$, which is given on generators by pull-back of structures. f^* only depends on the smooth homotopy class of f and is compatible with the $L(M)$ -module structures.

We define natural maps $H^{\text{odd}}(M, \mathbb{R}) \rightarrow \bar{L}(M)$ and $\bar{L}(M) \rightarrow L(M)$ by $[\rho] \mapsto [0, 0, 0, \rho]$ and $[\mathcal{F}, q, J, \rho] \mapsto [\mathcal{F}, q]$. Then we have the following exact sequence of $L(M)$ -modules

$$H^{\text{odd}}(M, \mathbb{R}) \rightarrow \bar{L}(M) \rightarrow L(M) \rightarrow H^{\text{ev}}(M, \mathbb{R})$$

(see [9], Prop. 21, for a similar argument).

Proposition 2.1 *The map $H^{\text{odd}}(M, \mathbb{R}) \rightarrow \bar{L}(M)$ is injective.*

Indeed, let $\omega \in \Omega^{\text{odd}}(M)$ be a closed form. If $(0, 0, 0, \omega) \sim 0$ in $\bar{L}_\epsilon(M)$, there there exists $(\mathcal{F}, q, J, \rho)$ together with two lagrangian subsheaves $\mathcal{L}_0, \mathcal{L}_1$ such that

$$[\omega] = [\tilde{\mathbf{ch}}(\nabla^{\oplus_0}, \nabla^{F_{\mathbb{C}}}) - \tilde{\mathbf{ch}}(\nabla^{\oplus_1}, \nabla^{F_{\mathbb{C}}})] = [\tilde{\mathbf{ch}}(\nabla^{\oplus_0}, \nabla^{\oplus_1})]$$

in $H^{odd}(M)$, where $\nabla^{\oplus i}$, $i = 0, 1$ are defined using the decompositions $F = L_i \oplus J(L_i)$. The right-hand side belongs to $\mathbf{ch}(K^{-1}(X))$ and is therefore rational. On the other hand it depends continuously on J . Therefore it is independent of J . First reducing to the case that $L_0 \cap L_1 = \{0\}$ and then choosing J such that $JL_i = L_{1-i}$, $i = 0, 1$, we see that $[\omega] = 0$.

We now construct a natural morphism of $L(M)$ -modules $\bar{\gamma} : \bar{L}(M) \rightarrow K_{\mathbb{R}/\mathbb{Z}}^{-1}(M)$. Here we use the definition of $K_{\mathbb{R}/\mathbb{Z}}^{-1}(M)$ in terms of generators and relations given in [8], Def. 5 + 6. A generator of $K_{\mathbb{R}/\mathbb{Z}}^{-1}(M)$ is a tuple (E, h^E, ∇^E, ρ) , where E is a $\mathbb{Z}_2$ -graded complex vector bundle of virtual dimension zero, h^E is a hermitean metric and ∇^E is a metric connection, both being compatible with the grading, and $\rho \in \Omega^{odd}(M)/\text{im}(d)$ satisfies $d\rho = \mathbf{ch}(\nabla^E)$. The relations of $K_{\mathbb{R}/\mathbb{Z}}^{-1}(M)$ are generated by

1. **isomorphism** $(E, h^E, \nabla^E, \rho) \sim (E', h^{E'}, \nabla^{E'}, \rho)$ if there exists an isomorphism from E to E' which is compatible with metrics and connections.
2. **direct sum** $(E, h^E, \nabla^E, \rho) + (E', h^{E'}, \nabla^{E'}, \rho') = (E \oplus E', h^{E \oplus E'}, \nabla^{E \oplus E'}, \rho + \rho')$
3. **change of connections** $(E, h^E, \nabla, \rho) \sim (E, h^E, \nabla', \rho')$ if $\rho' = \rho + \tilde{\mathbf{ch}}(\nabla', \nabla)$.
4. **trivial elements** If (E, h^E, ∇^E) is a $\mathbb{Z}_2$ -graded hermitean vector bundle with connection, then $(E \oplus E^{op}, h^{E \oplus E}, \nabla^{E \oplus E}, 0) \sim 0$, where E^{op} denotes E with the opposite grading.

Let $[E, h^E, \nabla^E, \rho]$ denote the class of (E, h^E, ∇^E, ρ) in $K_{\mathbb{R}/\mathbb{Z}}^{-1}(M)$.

We define $\bar{L}(M) \rightarrow K_{\mathbb{R}/\mathbb{Z}}^{-1}(M)$ by $\bar{\gamma}[\mathcal{F}, q, J, \rho] = [F_{\mathbb{C}}, h^{F_{\mathbb{C}}}, \nabla^{F_{\mathbb{C}}}, J, \rho]$, where $h^{F_{\mathbb{C}}}$ is the hermitean extension of the metric $J^* \circ Q$ on F .

The following diagram commutes

$$\begin{array}{ccc} \bar{L}(M) & \rightarrow & L(M) \\ \downarrow & & \downarrow \\ K_{\mathbb{R}/\mathbb{Z}}^{-1}(M) & \rightarrow & K^0(M) \end{array} .$$

We therefore obtain a natural map $\bar{L}(M) \rightarrow \bar{L}^{\mathbb{R}/\mathbb{Z}}(M)$ which is in fact surjective.

We now extend the functor $\bar{L}$ to $\mathcal{T}op_{smooth}$ by setting

$$\bar{L}(X) := \varprojlim_{f: M \rightarrow X} \bar{L}(M) ,$$

where the limit is taken over the category of manifolds over X . This extension has all functorial properties discussed above.

3 Eta invariants and index maps

3.1 The η -invariant

Using the η -invariant of the twisted signature operator for a closed odd-dimensional oriented manifold M in [9] Lott constructed a group homomorphism $\eta^{Lott} : \bar{L}^{Lott}(M) \rightarrow \mathbb{R}$. Its reduction modulo $\mathbb{Z}$ factors over the homomorphism $\eta^{\mathbb{R}/\mathbb{Z}} : K_{\mathbb{R}/\mathbb{Z}}^{-1}(M) \rightarrow \mathbb{R}/\mathbb{Z}$ constructed in [8], which is given by the pairing with the K -homology class induced by the odd signature operator.

Unfortunately, the homomorphism η^{Lott} does not factor over the quotient $\bar{L}^{Lott}(M) \rightarrow \bar{L}(M)$. In order to fix this for a given closed odd-dimensional oriented manifold M we define an extension

$$0 \rightarrow \mathbb{Z} \rightarrow \bar{L}_\epsilon^{ex}(M) \rightarrow L_\epsilon(M) \rightarrow 0 ,$$

such that

$$\bar{\eta} : \bar{L}_\epsilon^{ex}(M) \rightarrow \bar{L}_\epsilon(M) \rightarrow K_{\mathbb{R}/\mathbb{Z}}^{-1}(M) \xrightarrow{\eta^{\mathbb{R}/\mathbb{Z}}} \mathbb{R}/\mathbb{Z}$$

lifts to $\eta : \bar{L}_\epsilon^{ex}(M) \rightarrow \mathbb{R}$. Note that $\bar{L}_\epsilon^{ex}(M)$ is not a functor on M . Here $n := \dim(M)$, $\epsilon_n := [\frac{n(n-1)}{2}] \in \mathbb{Z}_2$, and we assume that $\epsilon = 1 - \epsilon_n$.

Let $(\mathcal{F}, q)$ be a generator of $L_\epsilon(M)$ and $\mathcal{L}$ be a lagrangian locally constant subsheaf of $\mathcal{F}$. We define an integer $\tau(\mathcal{F}, q, \mathcal{L})$ by the following construction. We consider the complex of sheaves

$$\mathcal{K}^\bullet : \mathcal{L} \xrightarrow{i} \mathcal{F} \xrightarrow{i^* \circ q} \mathcal{L}^* .$$

Let $(E_r^{\bullet\bullet}, d_r)$, $r \geq 1$ be the associated hyper cohomology spectral sequence. The duality q and the pairing between $\mathcal{L}$ and $\mathcal{L}^*$ induce a duality $q_{\mathcal{K}} : \mathcal{K} \xrightarrow{\sim} \mathcal{K}^*[-2]$ (the argument " $[-2]$ " indicates that $q_{\mathcal{K}}$ is a map of degree -2). The orientation of M and q induce a duality $q_{E_r} : E_r \xrightarrow{\sim} E_r^*[-2, -\dim(M)]$ (i.e. of bidegree $(-2, -\dim(M))$). Let N_r denote the total grading on E_r . Then we define the symmetric form Q_r on E_r by $Q_r(v, w) = q_{E_r}((-1)^{\frac{N_r(N_r-1)}{2}}v)(d_rw)$. The integer $\tau(\mathcal{F}, q, \mathcal{L})$ is now given by

$$\tau(\mathcal{F}, q, \mathcal{L}) := 2(\text{sign}(Q_1) + \text{sign}(Q_2)) .$$

We first define an extension $L_\epsilon^{ex}(M)$ of $L_\epsilon(M)$. A generator of the semigroup group $\hat{L}_\epsilon^{ex}(M)$ is a triple $(\mathcal{F}, q, z)$ consisting of a locally constant sheaf of finite-dimensional $\mathbb{R}$ -modules, an ϵ -symmetric duality structure q , and an integer z . The semigroup operation is given by

$$(\mathcal{F}, q, z) + (\mathcal{F}', q', z') := (\mathcal{F} \oplus \mathcal{F}', q \oplus q', z + z') .$$

Then we form $L_\epsilon^{ex}(M)$ by introducing the relation **lagrangian reduction**

$$(\mathcal{F}, q, z) \sim (0, 0, z + \tau(\mathcal{F}, q, \mathcal{L}))$$

if $\mathcal{F}$ admits a locally constant lagrangian subsheaf $\mathcal{L}$.

There is an exact sequence

$$0 \rightarrow \mathbb{Z} \rightarrow L_\epsilon^{ex}(M) \rightarrow L_\epsilon(M) \rightarrow 0 ,$$

where the maps are the obvious inclusion and projection. We now define $\bar{L}_\epsilon^{ex}(M)$ by the following pull-back diagram:

$$\begin{array}{ccc} \bar{L}_\epsilon^{ex}(M) & \rightarrow & L_\epsilon^{ex}(M) \\ \downarrow & & \downarrow \\ \bar{L}_\epsilon(M) & \rightarrow & L_\epsilon(M) \end{array} .$$

Then we have exact sequences

$$0 \rightarrow \mathbb{Z} \rightarrow \bar{L}_\epsilon^{ex}(M) \rightarrow \bar{L}_\epsilon(M) \rightarrow 0$$

and

$$H^{4*-(-1)^\epsilon}(M, \mathbb{R}) \rightarrow \bar{L}_\epsilon^{ex}(M) \rightarrow L_\epsilon^{ex}(M) \rightarrow H^{4*+1-(-1)^\epsilon}(M, \mathbb{R})$$

in a natural way.

An element of $\bar{L}_\epsilon^{ex}(M)$ can be represented by a tuple $(\mathcal{F}, q, J^F, \rho, z)$. We obtain a class $[\mathcal{F}, q, J^F, \rho] \in \bar{L}_\epsilon^{Lott}(M)$. If we set

$$\eta(\mathcal{F}, q, J^F, \rho, z) := \eta^{Lott}([\mathcal{F}, q, J^F, \rho]) - z ,$$

then we have the following result.

Theorem 3.1 *The map $(\mathcal{F}, q, J^F, \rho, z) \mapsto \eta(\mathcal{F}, q, J^F, \rho, z)$ induces a well-defined homomorphism $\eta : \bar{L}_\epsilon^{ex}(M) \rightarrow \mathbb{R}$ which lifts $\bar{\eta}$.*

We must show that η is well-defined with respect to lagrangian reduction. The idea is to consider the η -invariant of the signature operator on M twisted with the complex $K = \text{bundle}(\mathcal{K})$ and to rescale its differential. The corresponding adiabatic limits can be understood by the methods developed in [3] and [6] without investing essentially new ideas.

3.2 The secondary index map

Let $M \rightarrow B$ be a smooth locally trivial fibre bundle with even-dimensional closed fibres over a compact base B such that the vertical bundle TM/B is oriented. We set $n := \dim(TM/B)$.

In [9] Lott constructed a secondary index map $\pi_*^{\bar{L}, Lott} : \bar{L}_\epsilon^{Lott}(M) \rightarrow L_{\epsilon+\epsilon_n}^{Lott}(B)$ which fits into the commutative diagram

$$\begin{array}{ccccc} H^{4*-(-1)^\epsilon}(M, \mathbb{R}) & \rightarrow & \bar{L}_\epsilon^{Lott}(M) & \rightarrow & \bar{L}_\epsilon^{\mathbb{R}/\mathbb{Z}}(M) \\ \pi_*^{sign} \downarrow & & \pi_*^{\bar{L}, Lott} \downarrow & & \pi_*^{\bar{L}, \mathbb{R}/\mathbb{Z}} \downarrow \\ H^{4*-(-1)^{\epsilon+\epsilon_n}}(M, \mathbb{R}) & \rightarrow & \bar{L}_{\epsilon+\epsilon_n}^{Lott}(B) & \rightarrow & \bar{L}_{\epsilon+\epsilon_n}^{\mathbb{R}/\mathbb{Z}}(B) \end{array} .$$

Theorem 3.2 *By passing to quotients the map $\pi_*^{\bar{L}, Lott}$ induces a well-defined secondary index map*

$$\pi_*^{\bar{L}} : \bar{L}_\epsilon(M) \rightarrow \bar{L}_{\epsilon+\epsilon_n}(B) .$$

The idea of the proof consists in investigating the adiabatic limits of the η -form of the fibrewise signature operator twisted with the complex K under rescaling the differential. The arguments are similar to the case of analytic torsion forms. It suffices to adapt the methods developed in [4], [1], [2], [10], [11], [12], [13].

The following diagram commutes

$$\begin{array}{ccccc} H^{4*-(-1)^\epsilon}(M, \mathbb{R}) & \rightarrow & \bar{L}_\epsilon(M) & \rightarrow & L_\epsilon(M) \\ \pi_*^{sign} \downarrow & & \pi_*^{\bar{L}} \downarrow & & \pi_*^L \downarrow \\ H^{4*-(-1)^{\epsilon+\epsilon_n}}(M, \mathbb{R}) & \rightarrow & \bar{L}_{\epsilon+\epsilon_n}(B) & \rightarrow & L_{\epsilon+\epsilon_n}(B) \end{array} .$$

The secondary index map is natural with respect to pull-back of fibre bundles. Furthermore, if $\pi_1 : X \rightarrow X_1$ and $\pi_2 : X_2 \rightarrow B$ is an iterated bundle with oriented vertical bundles, and $\pi = \pi_2 \circ \pi_1$, then

$$\textbf{Theorem 3.3} \quad \pi_*^{\bar{L}} = (\pi_2^{\bar{L}})_* \circ (\pi_1^{\bar{L}})_*$$

This follows from the functoriality of the η -form studied in [11] in a similar way as the functoriality of the secondary index in [5] was deduced from the functoriality of the higher analytic torsion form [10].

3.3 The index map for L^{ex} and $\bar{L}^{ex}$

Let $\pi : M \rightarrow B$ as above and assume that B is oriented, closed, of dimension m . Furthermore, assume that $\epsilon = 1 - \epsilon_n - \epsilon_m$. Let $(\mathcal{F}, q)$ be a generator of $L_\epsilon(M)$. We define an integer $\tau(\mathcal{F}, q, M \xrightarrow{\pi} B)$ by the following construction. Let $(E^{\bullet\bullet}, d_r)$ be the Leray-Serre spectral sequence converging to sheaf cohomology $H^*(M, \mathcal{F})$ with $E_2^{p,q} = H^p(B, R^q \pi_* \mathcal{F})$. The orientations of TM/B and B and the duality q induce dualities $q_{E_r} : E_r \rightarrow E_r^*[-m, -n]$ which give the Poincaré duality on the limit. We define the symmetric form Q_r on E_r by $Q_r(v, w) = q_{E_r}((-1)^{\frac{N_r(N_r-1)}{2}} v)(d_r w)$, where N_r denotes the total degree. Then we set

$$\tau(\mathcal{F}, q, M \xrightarrow{\pi} B) := 2 \sum_{r \geq 2} \text{sign}(Q_r) .$$

Theorem 3.4 *1. The prescription*

$$\pi_*^{L^{ex}}[\mathcal{F}, q, z] := [H^\bullet R\pi_* \mathcal{F}, \pi_*(q), z - \tau(\mathcal{F}, q, M \xrightarrow{\pi} B)] .$$

defines an extended index map $\pi_^{L^{ex}} : L_\epsilon^{ex}(M) \rightarrow L_{\epsilon+\epsilon_n}^{ex}(B)$.*

2. *The extended index map is functorial with respect to iterated fibre bundles.*

The following diagram

$$\begin{array}{ccccc} \bar{L}_\epsilon(M) & \rightarrow & L_\epsilon(M) & \leftarrow & L_\epsilon^{ex}(M) \\ \pi_*^{\bar{L}} \downarrow & & \pi_*^L \downarrow & & \pi_*^{L^{ex}} \downarrow \\ \bar{L}_{\epsilon+\epsilon_n}(B) & \rightarrow & L_{\epsilon+\epsilon_n}(B) & \leftarrow & L_{\epsilon+\epsilon_n}^{ex}(B) \end{array}$$

commutes and induces an extended secondary index map $\pi_*^{\bar{L}^{ex}} : \bar{L}^{ex}(M)_\epsilon \rightarrow \bar{L}_{\epsilon+\epsilon_n}^{ex}(B)$ which is functorial with respect to iterated fibre bundles. We have the following compatibility of this extended secondary index map with the η -homomorphism.

Theorem 3.5

$$\begin{array}{ccc} \bar{L}_\epsilon^{ex}(M) & \xrightarrow{\eta} & \mathbb{R} \\ \pi_*^{\bar{L}^{ex}} \downarrow & & \parallel \\ \bar{L}_{\epsilon+\epsilon_n}^{ex}(B) & \xrightarrow{\eta} & \mathbb{R} \end{array}$$

This theorem essentially follows from Dai's adiabatic limit formula for the η -invariant [6].

References

- [1] A. Berthomieu and J.M. Bismut. Quillen metrics and higher analytic torsion forms. *J. Reine Angew. Math.*, 457(1994), 85–184.
- [2] J. M. Bismut. Families of immersions, and higher analytic torsion. *Asterisque*, 244, 1997.
- [3] J. M. Bismut and J. Cheeger. η -invariants and their adiabatic limits. *J. AMS*, 2(1989), 33–70.
- [4] J.M. Bismut and G. Lebeau. Complex immersions and Quillen metrics. *Publ. Math. IHES*, 74, 1991.
- [5] U. Bunke. On the functoriality of Lott's secondary analytic index. Preprint, 2000.
- [6] X. Dai. Adiabatic limits, non-multiplicativity of signature and Leray spectral sequence. *J. A.M.S.*, 4(1991), 265–321.
- [7] M. Karoubi. Homologie cyclique et K-théorie. *Asterisque*, 149, 1987.
- [8] J. Lott. $\mathbf{R}/\mathbf{Z}$ -index theory. *Comm. Anal. Geom.*, 2(1994), 279–311.
- [9] J. Lott. Secondary analytic indices. *Preprint*, 1999.

- [10] X. Ma. Functoriality of real analytic torsion form. Preprint, 1999.
- [11] X. Ma and U. Bunke. Flat duality bundle and the eta form. Preprint, 2000.
- [12] X. Ma. Formes de torsion analytique et familles de submersions I. *Bull. Soc. Math. France*, 127(1999), 541–621.
- [13] X. Ma. Formes de torsion analytique et familles de submersions II. *Asian J. of Math.*, 4(2000), 633-668.